

Restriction
Endonuclease



FauND I



Recognition
Sequence:

CA↓TATG
GTAT↑AC

S

E009T

1,000 units
20,000 u/ml

Lot: 41

Exp: 05/19

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	75-100	10-25	50-75	100	100

37°C

65°C

Y

λ

RE

TURBO

CERTIFICATE OF ANALYSIS

Description: Turbo FauND I can be used for short time (10-15 min) DNA digestion as well as for standard reaction. The reaction can be performed using optimal or universal (ROSE) Buffer. Buffer ROSE is perfect for double digestion.

Source: An *E.coli* strain that carries the cloned FauND I gene from *Flavobacterium aquatile ND*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1x SE-Buffer Y or 1x SE-Buffer ROSE. Incubate at 37°C

1X SE-Buffer Y (pH 7.9@ 25°C): 33 mM Tris-Ac, 66 mM KAc, 10 mM MgAc, 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65°C for 20 min.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with FauND I, approximately 80% of the DNA fragments can be ligated with high-activity T4 DNA Ligase and recut. In the presence of 10 % PEG ligation is better.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Reagents Supplied with Enzyme:

10x SE-Buffer Y, 10x SE-Buffer ROSE

Turbo DNA Digestion:

Applications:

- Fast DNA analysis
- Fast preparation of vectors for cloning
- Double digestion

Enzyme Properties:

1 µl of Turbo FauND I cuts 1 µg of DNA in 1x SE-Buffer Y or universal 1x SE-Buffer ROSE in 10-15 min (see the protocol below). Short time DNA digestion requires high quality purification of DNA sample. This enzyme can digest DNA at standard incubation time (1-16 hours) as well.

Turbo reaction protocol:

20 µl of the reaction volume:

- Reaction Buffer (x10) - 2 µl
- Plasmid DNA - 1-2 µl (up to 1 µg) or
- PCR product - 5-10 µl (~0,2 µg)
- Sterile water - up to 20 µl

+ 1 µl of Turbo Restriction Endonuclease
Incubate at 37°C for 10-15 min

For more details
scan the code



Ph/F +7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com