

Restriction  
Endonuclease



# Pst I



Recognition  
Sequence:

CTGCA↓G  
G↑ACGTC

**S** **E109T**  
4,000 units  
20,000 u/ml

Lot: 123  
Exp: 05/20  
Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	10-25	25-50	100	25-50	25-50	50

37°C No O λ RE TURBO

## CERTIFICATE OF ANALYSIS

**Description:** Turbo Pst I can be used for short time (10-15 min) DNA digestion as well as for standard reaction. The reaction can be performed using optimal or universal (ROSE) Buffer. Buffer ROSE is perfect for double digestion.

**Source:** An *E.coli* strain that carries the cloned Pst I gene from *Providencia stuartii*.

**Supplied in:**  
10 mM Tris-HCl (pH 7.6), 200 mM NaCl, 0.1 mM EDTA, 200 µg/ml BSA, 7 mM 2-mercaptoethanol, 50% glycerol.

**Reaction Conditions:**  
1x SE-Buffer O or 1x SE-Buffer ROSE. Incubate at 37°C  
1X SE-Buffer O (pH 7.6@ 25°C): 50 mM Tris-HCl, 100 mM NaCl, 10 mM MgCl<sub>2</sub>, 1 mM DTT

**Heat Inactivation:**  
Enzyme is not inactivated by incubation at 65° C for 20 minutes.

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

### Quality Control Assays

**Ligation:** After 20-fold overdigestion with Pst I, approximately 90% of the DNA fragments can be ligated with high-activity T4 DNA Ligase and recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 20 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

**Oligonucleotide Assay:** No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

**Reagents Supplied with Enzyme:**  
10x SE-Buffer O, 10x SE-Buffer ROSE

### Turbo DNA Digestion:

**Applications:**  
-Fast DNA analysis  
-Fast preparation of vectors for cloning  
-Double digestion

### Enzyme Properties:

1 µl of Turbo Pst I cuts 1 µg of DNA in 1x SE-Buffer O or universal 1x SE-Buffer ROSE in 10-15 min (see the protocol below). Short time DNA digestion requires high quality purification of DNA sample. This enzyme can digest DNA at standard incubation time (1-16 hours) as well.

### Turbo reaction protocol:

20 µl of the reaction volume:  
Reaction Buffer (x10) - 2 µl  
Plasmid DNA - 1-2 µl (up to 1 µg) or  
PCR product - 5-10 µl (~0,2 µg)  
Sterile water - up to 20 µl  
+ 1 µl of Turbo Restriction Endonuclease  
Incubate at 37°C for 10-15 min

For more details  
scan the code



Ph/F +7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com