

Restriction  
Endonuclease



# Ahl I



Recognition  
Sequence:

A↓CTAGT  
TGATC↑A

**S** **E173T**  
1,000 units  
20,000 u/ml

Lot: 15  
Exp: 09/20  
Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	75-100	25-50	25-50	75-100	100

37°C No B T7 TURBO

## CERTIFICATE OF ANALYSIS

Description: Turbo Ahl I can be used for short time (10-15 min) DNA digestion as well as for standard reaction. The reaction can be performed using optimal or universal (ROSE) Buffer. Buffer ROSE is perfect for double digestion.

Source: *Alteromonas haloplanktis SP*

Supplied in: 10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1 x SE-Buffer B or 1 x SE-Buffer ROSE.  
Incubate at 37°C.

1 x SE-Buffer B (pH 7.6@ 25°):

10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>                      1 mM DTT

Heat Inactivation: No (80°C for 20 minutes).

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of T7 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 20-fold overdigestion with Ahl I, >90% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 40 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Reagents Supplied with Enzyme:

10 x SE-Buffer B, 10 x SE-Buffer ROSE.

Turbo DNA Digestion:

Applications:

- Fast DNA analysis
- Fast preparation of vectors for cloning
- Double digestion

Enzyme Properties:

1 µl of Turbo Ahl I cuts 1 µg of DNA in 1 x SE-Buffer B or universal 1 x SE-Buffer ROSE in 10-15 min (see the protocol below). Short time DNA digestion requires high quality purification of DNA sample. This enzyme can digest DNA at standard incubation time (1-16 hours) as well.

Turbo reaction protocol:

20 µl of the reaction volume:

Reaction Buffer (x10) - 2 µl  
Plasmid DNA - 1-2 µl (up to 1 µg) or  
PCR product - 5-10 µl (~0,2 µg)  
Sterile water - up to 20 µl

+ 1 µl of Turbo Restriction Endonuclease  
Incubate at 37°C for 10-15 min.

For more details  
scan the code



Ph/F +7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com