

Restriction  
Endonuclease



# BamH I



Recognition  
Sequence:

G↓GATCC  
CCTAG↑G

Lot: 111  
Exp: 09/20  
Store at -20C

**E022T**  
20,000 units  
20,000 u/ml

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	25-50	100	75-100	75-100	25-50	100

37°C 65°C G λ RE TURBO

For more details  
scan the code



Ph/F +7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

**Description:** Turbo BamHI can be used for short time (10-15 min) DNA digestion as well as for standard reaction. The reaction can be performed using optimal or universal (ROSE) Buffer. Buffer ROSE is perfect for double digestion.

**Source:** An *E.coli* strain that carries the cloned BamHI gene from *Bacillus amyloliquefaciens* H.

**Supplied in:**  
50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 100 µg/ml BSA, 50% glycerol.

**Reaction Conditions:**  
1x SE-Buffer G or 1x SE-Buffer ROSE. Incubate at 37°C  
1X SE-Buffer G (pH 7.6@ 25°C): 10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 50 mM NaCl, 1 mM DTT

**Heat Inactivation:**  
Enzyme is inactivated by incubation at 65°C for 20 min

**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

### Quality Control Assays

**Ligation:** After 50-fold overdigestion with BamH I, approximately 90% of the DNA fragments can be ligated with high-activity T4 DNA Ligase and recut.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of DNA and 20 units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

**Oligonucleotide Assay:** No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

**Reagents Supplied with Enzyme:**  
10x SE-Buffer G, 10x SE-Buffer ROSE

### Turbo DNA Digestion:

#### Applications:

- Fast DNA analysis
- Fast preparation of vectors for cloning
- Double digestion

#### Enzyme Properties:

1 µl of Turbo BamHI cuts 1 µg of DNA in 1x SE-Buffer G or universal 1x SE-Buffer ROSE in 10-15 min (see the protocol below). Short time DNA digestion requires high quality purification of DNA sample. This enzyme can digest DNA at standard incubation time (1-16 hours) as well.

#### Turbo reaction protocol:

20 µl of the reaction volume:  
Reaction Buffer (x10) - 2 µl  
DNA (including plasmid) - 1-2 µl (up to 1 µg) or  
PCR product (purified) - 2-5 µl (~0.2 µg)  
Sterile water - up to 20 µl  
+ 1 µl of Turbo Restriction Endonuclease  
Incubate at 37°C for 10-15 min