

Restriction  
Endonuclease



# Acc36 I

Recognition  
Sequence:

ACCTGC(N)<sub>4</sub> ↓  
TGGACG(N)<sub>8</sub> ↑

**S**

**E289**

100 units  
2,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	25-50	25-50	50-75	50-75	100	100

37°C

65°C

Y

λ

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Acinetobacter calcoaceticus* 36.

### Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%  
glycerol.

### Reaction Conditions:

1x SE-Buffer Y. Incubate at 37° C.

### 1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac    66 mM KAc  
10 mM MgAc    1 mM DTT

### Heat Inactivation:

Enzyme is inactivated by incubation at 65 °C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of λ DNA in 1 hour at  
37° C in a total reaction volume of 50 µl.

### Quality Control Assays

Ligation: After 3-fold overdigestion with Acc36 I, 90%  
of the DNA fragments can be ligated with T4 DNA Ligase  
and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of  
DNA and 4 Units of enzyme incubated for 16 hours resulted  
in the same pattern of DNA bands as a reaction incubated  
for 1 hour.

High enzyme concentration may result in star activity.

Oligonucleotide Assay: No detectable degradation of  
a single-stranded and double-stranded oligonucleotide  
was observed after incubation with 2 units of restriction  
endonuclease for 3 hours.

### Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

### Reagents Supplied with Enzyme:

10X SE Buffer Y.

### Note:

The single Acc36I sites in pBR322 and pUC19 are  
resistant to cleavage.