

Restriction  
Endonuclease



## AccBS I

Recognition  
Sequence:

CCG↓CTC  
GGC↑GAG

XS

**E007m**

250 units  
5,000 u/ml

Lot:

Exp:

**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	75-100	25-50	25-50	100	100

37°C

65°C

λ

minimal

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Acinetobacter calcoaceticus* BS.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA,  
50% glycerol.

Reaction Conditions:

1x SE-Buffer Y. Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac    66 mM KAc  
10 mM MgAc    1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65 °C for 20  
minutes.

Unit Definition: One unit of the enzyme is the amount  
required to hydrolyze 1 µg of λ DNA in 1 hour at 37° C  
in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with AccBS I,  
90% of the DNA fragments can be ligated with T4  
DNA Ligase and 50% of these can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg  
of DNA and 10 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of  
a single-stranded and double-stranded oligonucleotide  
was observed after incubation with 5 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE-Buffer Y.