

Restriction
Endonuclease



Bpu14 I

Recognition
Sequence:

TT↓CGAA
AAGC↑TT

S

E033

1000 units
10,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	100	25-50	25-50	75-100	100

37°C

65°C

G

λ

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Bacillus pumilus* 14.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

1× SE-Buffer G. Incubate at 37° C.

1X SE-Buffer G (pH 8.5 @ 25° C):

10 mM Tris-Ac 50 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65 °C for 20
minutes.

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of Lambda DNA in 1
hour at 37°C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Bpu14 I,
95% of the DNA fragments can be ligated with T4 DNA
Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg
of DNA and 20 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a reaction
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of
a single-stranded and double-stranded oligonucleotide
was observed after incubation with 10 units of restriction
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer G