

Restriction
Endonuclease



AccB7 I

Recognition
Sequence: CCANNNN↓NTGG
GGTN↑NNNNACC

S

E179

200 units
5,000 u/ml

Lot:

Exp:

Store at -20°C

| SE-Buffers | B | G | O | W | Y | ROSE |
|------------|-------|-----|-------|-------|-------|------|
| %Activity | 10-25 | 100 | 25-50 | 50-75 | 50-75 | 100 |

37°C

65°C

G

λ

Dcm

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Acinetobacter calcoaceticus* B7.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

1x SE-Buffer G. Incubate at 37° C.

1X SE-Buffer G (pH 7.6 @ 25° C):

10 mM Tris-HCl 50 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65 °C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA (Dcm-) in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with AccB7 I, 95% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 5 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Star activity: High enzyme concentration may result in star activity.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer G.

Blocked by overlapping Dcm methylation (C^mCWGG):
CCANNNCCTGG or CCAGGNNNTGG.