

Restriction
Endonuclease



Ahl I

Recognition
Sequence:

A↓CTAGT
TGATC↑A

XS

E173m
250 units
10,000 u/ml

Lot:
Exp:
Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	75-100	25-50	25-50	75-100	100

37°C **No** **B** **T7** **minimal**

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Alteromonas haloplanktis* SP.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM KCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%
glycerol.

Reaction Conditions:

1x SE-Buffer B, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

No (80° C for 20 minutes).

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of T7 DNA in 1 hour
at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Ahl I, >90%
of the DNA fragments can be ligated with T4 DNA Ligase
and recut.

16-Hour Incubation: A 50 µl reaction containing in 1 µg
of DNA and 20 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a reaction
incubated for 1 hour.
No using BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a
single-stranded and double-stranded oligonucleotide
was observed after incubation with 10 units of restriction
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer B, BSA (10 mg/ml).