

Restriction  
Endonuclease



# Sfr303 I

Recognition  
Sequence:

CCGC↓GG  
GG↑CGCC

XS

**E127m**  
250 units  
5,000 u/ml

Lot:  
Exp:  
**Store at -20°C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	50-75	10-25	10-25	75-100	100

37°C

65°C

B

λ

minimal

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Streptomyces fradiae* 303.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:

1x SE-Buffer B. Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of Lambda DNA in 1  
hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with Sfr303 I, >  
90% of the DNA fragments can be ligated with T4 DNA  
Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing in 1 µg  
of DNA and 10 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 5 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer B.