

Restriction
Endonuclease



BspI720 I

Recognition
Sequence:

GC↓TNAGC
CGANT↑CG

S

E185

500 units
10,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	100	50-70	50-75	75-100	75

37°C

80°C

G

λ

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Bacillus species 1720*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

1x SE-Buffer G. Incubate at 37° C.

1X SE-Buffer G (pH 7.6 @ 25° C):

10 mM Tris-HCL 50 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80°C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Bsp1720 I, about 80% of the DNA fragments can be ligated with T4 DNA Ligase and 95% of these can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer G.