

Restriction
Endonuclease



Vsp I

Recognition
Sequence:

AT↓TAAT
TAAT↑TA

XS

E139m
500 units
10,000 u/ml

Lot:
Exp:
Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	10-25	50-75	100	25-50	50

37°C **65°C** W λ RR minimal

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *Vsp I* gene from *Vibrio species 343*.

Supplied in:
10 mM Tris-HCl (pH 7.6), 50 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:
1X SE-Buffer W. Incubate at 37° C.

1X SE-Buffer W (pH 8.5 @ 25° C):
10 mM Tris-HCl 100 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:
Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of lambda DNA in 1 hour at 65° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Vsp I, ~70% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer W.

Blocked by ATTA^mAT methylation.