

Restriction
Endonuclease



PciS I

Recognition
Sequence:

GCTCTTC(N)₁↓
CGAGAAG(N)₄↑

S

E497
50 units
500 u/ml

Lot:
Exp:
Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	50-75	0-10	0-10	75-100	50

37°C 65°C B λ

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Planococcus citreus S.*

Supplied in:
10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%
glycerol.

Reaction Conditions:
1X SE-Buffer B. Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):
33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:
Enzyme is inactivated by incubation at 65° C for 20
minutes.

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of Lambda DNA in
1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays
Ligation: After 3-fold overdigestion with PciS I, ~90%
of the DNA fragments can be ligated with T4 DNA
Ligase and ~95% of these can be recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg
of DNA and 0.5 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a reaction
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a
single-stranded and double-stranded oligonucleotide
was observed after incubation with 0.5 units of restriction
endonuclease for 3 hours.

Enzyme Properties:
When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer B.