

Restriction  
Endonuclease



# Pps I

Recognition  
Sequence:

GAGTC(N)<sub>4</sub> ↓  
CTCAG(N)<sub>5</sub> ↑

S

**E269**  
25 units  
500 u/ml

Lot:  
Exp:  
**Store at -20°C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	10-25	0-10	25-50	100	50

37°C

65°C

Y

λ

BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Pseudomonas pseudoalcaligenes*.

Supplied in:  
10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:  
1X SE-Buffer Y, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):  
33 mM Tris-Ac      66 mM KAc  
10 mM MgAc      1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 65° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of Lambda DNA in 1  
hour at 37° C in a total reaction volume of 50 µl.  
To obtain 100% activity, BSA should be added to the  
1x reaction mix to a final concentration of 100 µg/ml.

Quality Control Assays  
Ligation: After 2-fold overdigestion with Pps I, 20% of  
the DNA fragments can be ligated with T4 DNA Ligase  
and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of  
DNA and 1 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Do not use BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 0.5 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:  
When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:  
10X SE Buffer Y, BSA (10 mg/ml).