

Restriction
Endonuclease



PspL I

Recognition
Sequence:

C↓GTACG
GCATG↑C

S

E223

200 units
2,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	75-100	25-50	10-25	100	100

37°C

65°C

Y

λ/HindIII

BSE

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Pseudomonas species L.*

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%
glycerol.

Reaction Conditions:

1X SE-Buffer Y, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20
minutes.

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of Lambda DNA/
HindIII in 1 hour at 37° C in a total reaction volume
of 50 µl.

To obtain 100% activity, BSA should be added to
the 1x reaction mix to a final concentration of 100
µg/ml.

Quality Control Assays

Ligation: After 2-fold overdigestion with PspL I, >
95% of the DNA fragments can be ligated with T4
DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg
of DNA and 2 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a
reaction incubated for 1 hour.

Do not use BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of
a single-stranded and double-stranded oligonucleotide
was observed after incubation with 2 units of restriction
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y, BSA (10 mg/ml).