

Restriction
Endonuclease



TseF I

Recognition
Sequence:

↓GTSA
CASTG↑

S

E589

200 units
5,000 u/ml

Lot:

Exp:

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	50-75	0-10	25-50	50-75	50

65°C

No

B

λ

RR

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *TseFI* gene from *Thermus species F35*.

Supplied in:

10 mM Tris-HCl (pH 7.6), 50 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA, and 50% glycerol.

Reaction Conditions:

1X SE-Buffer B. Incubate at 65° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is not inactivated by incubation at 80° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 65° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with TseF I, more than 95% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 5 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer B.