

Restriction  
Endonuclease



# Hga I

Recognition  
Sequence:

GACGC(N)<sub>5</sub>↓  
CTGCC(N)<sub>10</sub>↑

S

**E461**

50 units  
1,000 u/ml

Lot:

Exp:

**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	75-100	10-25	25-50	50-75	50

37°C

65°C

B

pBR322

RR

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *Hga I* gene from *Haemophilus gallinarum*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 100 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1X SE-Buffer B. Incubate at 37°C.

1X SE-Buffer B (pH 7.6 @ 25°C):

10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65°C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 3-fold overdigestion with Hga I, more than 90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 0.5 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Incubation with > 2 units of HgaI per 1 µg of DNA and digestion > 1 hour are not recommended.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 1 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer B.

Blocked by CG methylation.