

Restriction  
Endonuclease



# Mox20 I

Recognition  
Sequence:

TGG↓CCA  
ACC↑GGT

S

E301

1,000 units  
20,000 u/ml

Lot:

Exp:

Store at -20°C

| SE-Buffers | B     | G     | O   | W      | Y     | ROSE |
|------------|-------|-------|-----|--------|-------|------|
| %Activity  | 10-25 | 25-50 | 100 | 75-100 | 25-50 | 75   |

37°C

No

O

λ

Dcm

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Microbacterium oxydans*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 200 mM KCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:

1X SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):

50 mM Tris-HCl      100 mM NaCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:

NO (80° C for 20 minutes).

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (Dcm-) in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 20-fold overdigestion with Mox20 I, 80% of the DNA fragments can be ligated and recut. More complete ligation can be reached by using 10% PEG.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 40 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer O.

Cleaved of DNA is impaired by overlapping Dcm methylation (C<sup>m</sup>CWGG): TGGCCAGG.