

Restriction
Endonuclease



Fok I

Recognition
Sequence:

GGATC(N)₉ ↓
CCTAC(N)₁₃ ↑

S

E247

100 units
1,000 u/ml

Lot:

Exp:

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	50-75	25-50	25-50	100	100

37°C

65°C

Y

λ

RRR

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned *Fok I* gene from *Flavobacterium okeanokoites*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1X SE-Buffer Y. Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl. Overdigestions of > 5 units of Fok I per 1 µg of DNA and incubations > 2 hours are not recommended.

Quality Control Assays

Ligation: After 2-fold overdigestion with Fbl I, ~95% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 1 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 1 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y.