

Restriction  
Endonuclease



# Bme18 I

Recognition  
Sequence:

G↓GWCC  
CCWG↑G

XS

**E029m**  
500 units  
10,000 u/ml

Lot:  
Exp:  
**Store at -20°C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	10-25	25-50	100	75-100	10-25	80

37°C 65°C O λ minimal

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Bacillus megaterium 18.*

Supplied in:  
10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:  
1X SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):  
50 mM Tris-HCl    100 mM NaCl  
10 mM MgCl<sub>2</sub>    1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 65° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of lambda DNA in 1  
hour at 37° C in a total reaction volume of 50 µl.

### Quality Control Assays

Ligation: After 10-fold overdigestion with Bme 18 I, more  
than 90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of  
DNA and 20 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 10 units of restriction  
endonuclease for 3 hours.

### Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:  
10X SE Buffer O.

Cleaved of DNA is impaired by overlapping dcm-  
methylation  
(C<sup>m</sup>CWGG): GGWCCWGG.