

Restriction
Endonuclease



Hpa I

Recognition
Sequence:

GTT↓ACC
CAA↑TTG

XS

E077m
250 units
5,000 u/ml

Lot:
Exp:
Store at -20C

| SE-Buffers | B | G | O | W | Y | ROSE |
|------------|------|-------|-------|-------|-----|------|
| %Activity | 0-10 | 50-75 | 10-25 | 25-50 | 100 | 25 |

37°C 65°C Y λ RR BSA

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned *Hpa I* gene from *Haemophilus parainfluenzae*.

Supplied in:
10 mM Tris-HCl (pH 7.6), 50 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:
1X SE-Buffer Y. Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):
33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:
Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with Hpa I, ~60% of the DNA fragments can be ligated and recut.

16-Hour Incubation: Long incubation is not recommended owing to occurrence of star activity. Star activity is observed at a great than 5-fold overdigestion of 1 µg substrate with Hpa I.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer Y.