

Restriction
Endonuclease



Bbv12 I

Recognition
Sequence:

GWGCW↓C
C↑WCGWG

S

E023

200 units
2,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	10-25	100	75-100	10-25	60

37°C

80°C

O

λ

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Bacillus brevis* 12.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%
glycerol.

Reaction Conditions:

1X SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):

50 mM Tris-HCl 100 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80° C for 20
minutes.

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of lambda DNA in 1
hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Bbv 12 I, more
than 90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of
DNA and 4 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a reaction
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a
single-stranded and double-stranded oligonucleotide
was observed after incubation with 2 units of restriction
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer O.