

Restriction  
Endonuclease



# Bse118 I

Recognition  
Sequence:

R↓CCGGY  
YGGCCT↑R

S

**E039**

200 units  
5,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	50-75	100	75-100	25-50	100

65°C

80°C

O

λ

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Bacillus stearothermophilus 118.*

Supplied in:

10 mM KH<sub>2</sub>PO<sub>4</sub> (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:

1X SE-Buffer O. Incubate at 65° C.

1X SE-Buffer O (pH 7.6 @ 25° C):

50 mM Tris-HCl      100 mM NaCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of Lambda DNA/  
HindIII in 1 hour at 65° C in a total reaction volume  
of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with Bse118 I, ~  
90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of  
DNA and 10 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 10 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer O.