

Restriction  
Endonuclease



# PspC I

Recognition  
Sequence:

CAC↓GTG  
GTG↑CAC

XS

**E475m**  
500 units  
20,000 u/ml

Lot:  
Exp:  
**Store at \*  
-20°C/-70°C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	50-75	0-10	0-10	50-75	5

37°C

65°C

B

λ

BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Pseudomonas species C.*

Supplied in:  
10 mM Tris-HCl (pH 7.5), 50 mM KCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:  
1X SE-Buffer B, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):  
10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 65° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of Lambda DNA in  
1 hour at 37° C in a total reaction volume of 50 µl.  
To obtain 100% activity, BSA should be added to the  
1 x reaction mix to a final concentration of 100 µg/ml.

### Quality Control Assays

Ligation: After 20-fold overdigestion with PspC I, >  
90% of the DNA fragments can be ligated with T4  
DNA Ligase and recut. Ligation is better in presence  
of 10% PEG.

16-Hour Incubation: A 50 µl reaction containing 1  
µg of DNA and 40 Units of enzyme incubated for 16  
hours resulted in the same pattern of DNA bands as  
a reaction incubated for 1 hour.  
Do not use BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 20 units of restriction  
endonuclease for 3 hours.

### Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:  
10X SE Buffer B, BSA (10 mg/ml).

**Storage at -70° C is recommended for periods longer  
than 30 days.**