

Restriction
Endonuclease



Ege I

Recognition
Sequence:

GGC↓GCC
CCG↑CGG

S

E243

500 units
5,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	75-100	10-25	50-75	75-100	100

37°C

65°C

B

λ/HindIII

BSA

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Enterobacter gergoviae*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

11× SE-Buffer B, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA/HindIII in 1 hour at 37° C in a total reaction volume of 50 µl. To obtain 100% activity, BSA should be added to the 1x reaction mix to a final concentration of 100 µg/ml.

Quality Control Assays

Ligation: After 5-fold overdigestion with Ege I, 70% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

NO using BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer B, BSA (10 mg/ml).