

Restriction  
Endonuclease



# EcoR I

Recognition  
Sequence:

G↓AATTC  
CTTAA↑G

S

**E057X**  
5,000 units  
50,000 u/ml

Lot:  
Exp:  
**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	75-100	75-100	75-100	50-75	50

37°C 65°C EcoRI λ RR BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *EcoR I* gene from *Escherichia coli*.

Supplied in:  
10 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 100 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:  
1× SE-Buffer EcoR I, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer EcoR I (pH 7.6 @ 25° C):  
10 mM Tris-HCl 50mM NaCl  
10 mM MgCl<sub>2</sub> 1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 65° C for 20  
minutes.

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of Lambda DNA in 1  
hour at 37° C in a total reaction volume of 50 µl.  
To obtain 100% activity, BSA should be added to the  
1 x reaction mix to a final concentration of 100 µg/ml.  
High enzyme concentration and using of nonoptimal  
buffer may result in star activity.  
Do not use BSA for long incubation.

### Quality Control Assays

Ligation: After 40-fold overdigestion with EcoR I, ~95%  
of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing in 1 µg  
of DNA and 40 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 20 units of restriction  
endonuclease for 3 hours.

### Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

### Reagents Supplied with Enzyme:

10X SE Buffer EcoR I, BSA (10 mg/ml).