

Restriction
Endonuclease



Bsa29 I

Recognition
Sequence:

AT↓CGAT
TAGC↑TA

XS

E205m
200 units
20,000 u/ml

Lot:
Exp:
Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	25-50	100	50-75	50-75	75-100	100

37°C **65°C** **G** λ **Dam** **BSA**

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: *Bacillus stearothermophilus 29.*

Supplied in:
10 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA,
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%
glycerol.

Reaction Conditions:
1XSE-Buffer G, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer G (pH 7.6 @ 25° C):
10 mM Tris-HCl 50 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:
Enzyme is inactivated by incubation at 65°C for 20
minutes.

Unit Definition: One unit is defined as the amount of
enzyme required to digest 1 µg of Lambda DNA in 1
hour at 37° C in a total reaction volume of 50 µl.
To obtain 100% activity, BSA should be added to the
1x reaction mix to a final concentration of 100 µg/ml.

Quality Control Assays

Ligation: After 20-fold overdigestion with Bsa29 I, >
90% of the DNA fragments can be ligated with T4 DNA
Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of
DNA and 40 Units of enzyme incubated for 16 hours
resulted in the same pattern of DNA bands as a reaction
incubated for 1 hour.

Do not use BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a
single-stranded and double-stranded oligonucleotide
was observed after incubation with 20 units of restriction
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)
SE-Buffer, it may be necessary to add more enzymes
to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer G, BSA (10mg/ml).

Blocked by overlapping Dam methylation (G^mATC):
GATCGATC.

Blocked by CG methylation.