

Restriction
Endonuclease



SfaN I

Recognition
Sequence:

GCATC(N)₅↓
CGTAG(N)₉↑

XS

E165m
100 units
10,000 u/ml

Lot:
Exp:
Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	10-25	25-50	100	75-100	0-10	25

37°C

80°C

O

λ

RR

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *SfaN I* gene from *Streptococcus faecalis N*.

Supplied in:
10 mM Tris-HCl (pH 7.5), 300 mM NaCl, 0.1 mM EDTA,
200 µg/ml BSA, 1 mM DTT, 50% glycerol.

Reaction Conditions:
1X SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):
50 mM Tris-HCl 100 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:
Enzyme is inactivated by incubation at 80° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays
Ligation: After 10-fold overdigestion with SfaN I, more than 95% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:
When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:
10X SE Buffer O.