

Restriction
Endonuclease



Mnl I

Recognition
Sequence:

CCTC(N)₇↓
GGAG(N)₆↑

XS

E481m
150 units
10,000 u/ml

Lot:
Exp:
Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	100	25-50	25-50	75-100	100

37°C 65°C G λ RR BSA

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned *Mnl I* gene from *Moraxella nonliquefaciens*.

Supplied in:

10 mM Tris-HCl (pH 7.6), 200 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1X SE-Buffer G, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer G (pH 7.6 @ 25° C):

10 mM Tris-HCl 50 mM NaCl
10 mM MgCl₂ 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl. To obtain 100% activity, BSA should be added to the 1x reaction mix to a final concentration of 100 µg/ml.

Quality Control Assays

Ligation: After 10-fold overdigestion with Mnl I, 50% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Do not use BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer G, BSA (10mg/ml).

Blocked by overlapping CG methylation: CCT^m CG.