

Restriction  
Endonuclease



# Ahl I

Recognition  
Sequence:

A↓CTAGT  
TGATC↑A

L

**E174**

5,000 units  
20,000 u/ml

Lot:

Exp:

**Store at -20°C**

| SE-Buffers | B   | G      | O     | W     | Y      | ROSE |
|------------|-----|--------|-------|-------|--------|------|
| %Activity  | 100 | 75-100 | 25-50 | 25-50 | 75-100 | 100  |

37°C

No

B

T7

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Alteromonas haloplanktis SP.*

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM KCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 200 µg/ml BSA, 50%  
glycerol.

Reaction Conditions:

1x SE-Buffer B, BSA (100 µg/ml). Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:

No (80° C for 20 minutes).

Unit Definition: One unit is defined as the amount of  
enzyme required to digest 1 µg of T7 DNA in 1 hour  
at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Ahl I, >90%  
of the DNA fragments can be ligated with T4 DNA Ligase  
and recut.

16-Hour Incubation: A 50 µl reaction containing in 1 µg  
of DNA and 20 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.  
No using BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a  
single-stranded and double-stranded oligonucleotide  
was observed after incubation with 10 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer B, BSA (10 mg/ml).