

Restriction  
Endonuclease



# Bsp19 I

Recognition  
Sequence:

C↓CATGG  
GGTACT↑C

S

**E925**

1,000 units  
20,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	100	0-15	25-50	100	100

37°C

80°C

Y

λ

BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Bacillus species 19*.

Supplied in:

10 mM Tris-HCl (pH 7.4), 200 mM NaCl, 0.1 mM EDTA,  
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1× SE-Buffer Y, BSA (100 µg/ml). Incubate at 37 °C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac

66 mM KAc

10 mM MgAc

1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80 °C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl. To obtain 100% activity, BSA should be added to the 1x reaction mix to a final concentration of 100 µg/ml.

Quality Control Assays

Ligation: After 20-fold overdigestion with Bsp19 I, 95% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 40 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

No using BSA for long incubation.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y, BSA (10 mg/ml).

Bsp19I cuts hemi methylated site 5'-(5mC) CATGG-3' / 3'-GGTACC-5' and doesn't cut methylated sites 5'-(5mC) CATGG-3' / 3'-GGTAC(5mC)-5' and 5'-(4mC) CATGG-3' / 3'-GGTAC(4mC)-5'.