

Restriction  
Endonuclease



# Mlu I

Recognition  
Sequence:

A↓CGCGT  
TGCGCT↑A

S

**E917**

1,000 units  
20,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	100	100	100	100	100

37°C

NO

Y

λ

BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Micrococcus luteus*.

Supplied in:

10 mM Tris-HCl (pH 7.4), 200 mM NaCl, 0.1 mM EDTA,  
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1X SE-Buffer Y, BSA (100 µg/ml) . Incubate at 37 °C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac      66 mM KAc  
10 mM MgAc      1 mM DTT

Heat Inactivation:

NO ( 80°C for 20 minutes ).

Unit Definition:One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation:After 20-fold overdigestion with MluI, more than 90% of the DNA fragments can be ligated and recut.

16-Hour Incubation:A 50 µl reaction containing 1 µg of DNA and 40 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay:No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y, BSA (10 mg/ml).