

Restriction  
Endonuclease



# Hae III

Recognition  
Sequence:

GG↓CC  
CC↑GG

**S**

**E067**

2,000 units  
10,000 u/ml

Lot:

Exp:

**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	100	25-50	50-75	50-75	100

37°C

80°C

G

λ

RR

minimal

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *E.coli* strain that carries the cloned *Hae III* gene from *Haemophilus aegyptius*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA,  
1 mM DTT, 200 µg/ml BSA, and 50% glycerol.

Reaction Conditions:

1X SE-Buffer G. Incubate at 37° C.

1X SE-Buffer G (pH 7.6 @ 25° C):

10 mM Tris-HCl    50 mM NaCl  
10 mM MgCl<sub>2</sub>    1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 20-fold overdigestion with Hae III, ~90% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer G.