

Restriction  
Endonuclease



# Pci I

Recognition  
Sequence:

A↓CATGT  
TGTACT↑A

S

E275

300 units

10,000 u/ml

Lot:

Exp:

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	50-75	75-100	100	75-100	50-75	50

37°C 65°C O T7 RR minimal

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E. coli* strain that carries the cloned *Pci I* gene from *Planococcus citreus* SE-F45.

Supplied in:

10 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA, 7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

1x SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):

50 mM Tris-HCl 100 mM NaCl  
10 mM MgCl<sub>2</sub> 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of T7 DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 10-fold overdigestion with Pci I, ~90% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 5 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer O.

Blocked by<sup>m</sup>ACATGT methylation.

Not blocked by AC<sup>m</sup>ATGT methylation.