

Restriction  
Endonuclease



## Dri I

Recognition  
Sequence:

GACNNN↓NNGTC  
CTGNN↑NNNCAG

S

**E193**

500 units  
10,000 u/ml

Lot:

Exp:

Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	75-100	75-100	10-25	10-25	100	40

37°C

65°C

Y

λ

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: *Deinococcus radiodurans* EA.

Supplied in:

10 mM Tris-HCl (pH 7.5), 250 mM NaCl, 0.1 mM EDTA,  
7 mM 2-mercaptoethanol, 50% glycerol.

Reaction Conditions:

1x SE-Buffer Y. Incubate at 37°C.

1X SE-Buffer Y (pH 7.9 @ 25°C):

33 mM Tris-Ac    66 mM KAc  
10 mM MgAc    1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65°C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 5-fold overdigestion with Dri I, ~5% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

In the presence of 10% PEG ligation is better.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y.