

Restriction  
Endonuclease



# Hinf I

Recognition  
Sequence:

G↓ANTC  
CTNA↑G

XS

**E075m**  
500 units  
10,000 u/ml

Lot:  
Exp:  
**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	25-50	75-100	100	75-100	75-100	100

37°C

80°C

O

λ

RR

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned *Hinf I* gene from *Haemophilus influenzae*.

Supplied in:  
10 mM Tris-HCl (pH 7.5), 50 mM NaCl, 0.1 mM EDTA,  
200 µg/ml BSA, 1 mM DTT, 50% glycerol.

Reaction Conditions:  
1X SE-Buffer O. Incubate at 37° C.

1X SE-Buffer O (pH 7.6 @ 25° C):  
50 mM Tris-HCl      100 mM NaCl  
10 mM MgCl<sub>2</sub>      1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 80° C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

### Quality Control Assays

Ligation: After 20-fold overdigestion with *Hinf I*, ~90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 40 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

### Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

### Reagents Supplied with Enzyme:

10X SE Buffer O.