

Restriction  
Endonuclease



# Hind III

Recognition  
Sequence:

A↓AGCTT  
TTCGA↑A

XS

**E903m**  
5,000 units  
20,000 u/ml

Lot:  
Exp:  
**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	100	0-10	NR	100	100

37°C 80°C Y λ RR BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned Hind III gene from *Haemophilus influenzae Rd*.

Supplied in:  
10 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.1 mM EDTA,  
200 µg/ml BSA, 1 mM DTT, 50% glycerol.

Reaction Conditions:  
1x SE-Buffer Y, BSA (100 µg/µl).  
Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):  
33 mM Tris-Ac 66 mM KAc  
10 mM MgAc 1 mM DTT

Heat Inactivation:  
Enzyme is inactivated by incubation at 80 °C for 20  
minutes.

Unit Definition: One unit is defined as the amount  
required to digest 1 µg of λ DNA in 1 hour at 37° C  
in a total reaction volume of 50 µl. To obtain 100%  
activity, BSA should be added to the 1X reaction  
mix to final concentration of 100 µg/µl.

Quality Control Assays  
Ligation: After 50-fold overdigestion with Hind III,  
approximately 90% of the DNA fragments can be  
ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg  
of DNA and 40 Units of enzyme incubated for 16 hours  
resulted in the same pattern of DNA bands as a reaction  
incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of  
a single-stranded and double-stranded oligonucleotide  
was observed after incubation with 20 units of restriction  
endonuclease for 3 hours.

Enzyme Properties:  
When using a buffer other than the optimal (Supplied)  
SE-Buffer, it may be necessary to add more enzymes  
to achieve complete digestion.

Reagents Supplied with Enzyme:  
10X SE Buffer Y, BSA (10 mg/ml).