

Restriction  
Endonuclease



# Hind III

Recognition  
Sequence:

A↓AGCTT  
TTCGA↑A

S

**E903**

10,000 units  
20,000 u/ml

Lot:

Exp:

**Store at -20C**

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	0-10	100	0-10	NR	100	100

37°C

80°C

Y

λ

RR

BSA

For more details  
scan the code



Ph/F+7(383)333-6853  
info@sibenzyme.com  
www.sibenzyme.com

## CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned Hind III gene from *Haemophilus influenzae Rd*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 200 mM NaCl, 0.1 mM EDTA, 200 µg/ml BSA, 1 mM DTT, 50% glycerol.

Reaction Conditions:

1x SE-Buffer Y, BSA (100 µg/µl).  
Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac    66 mM KAc  
10 mM MgAc    1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80 °C for 20 minutes.

Unit Definition: One unit is defined as the amount required to digest 1 µg of λ DNA in 1 hour at 37° C in a total reaction volume of 50 µl. To obtain 100% activity, BSA should be added to the 1X reaction mix to final concentration of 100 µg/µl.

Quality Control Assays

Ligation: After 50-fold overdigestion with Hind III, approximately 90% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y, BSA (10 mg/ml).