

Restriction
Endonuclease



Hpa II

Recognition
Sequence:

C↓CGG
GGC↑C

S

E161

500 units
10,000 u/ml

Lot:

Exp:

Store at -20C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	100	50-75	10-25	25-50	50-75	50

37°C

80°C

B

λ

RR

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned gene Hpa II from *Haemophilus parainfluenzae*.

Supplied in:

10 mM Tris-HCl (pH 7.5), 50 mM NaCl, 0.1 mM EDTA, 100 µg/ml BSA, 1 mM DTT, 50% glycerol.

Reaction Conditions:

1x SE-Buffer B. Incubate at 37° C.

1X SE-Buffer B (pH 7.6 @ 25° C):

10 mM Tris-HCl 10 mM MgCl₂
1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 80 °C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to hydrolyze 1 µg of λ DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation : After 10-fold overdigestion with Hpa II, approximately 95% of the DNA fragments can be ligated with T4 DNA Ligase and recut.

16-Hour Incubation: A 50 µl reaction containing in 1 µg of DNA and 10 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 10 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE-Buffer B.

Blocked by CG methylation.