

Restriction
Endonuclease



Not I

Recognition
Sequence:

GC↓GGCCGC
CGCCGG↑CG

XS

E911m
300 units
20,000 u/ml

Lot:
Exp:
Store at -20°C

SE-Buffers	B	G	O	W	Y	ROSE
%Activity	10-25	100	10-25	50-75	100	100

37°C

65°C

Y

pUC102

For more details
scan the code



Ph/F+7(383)333-6853
info@sibenzyme.com
www.sibenzyme.com

CERTIFICATE OF ANALYSIS

Source: An *E.coli* strain that carries the cloned *Not I* gene from *Nocardia otitidis-caviarum*.

Supplied in:

10 mM Tris-HCl (pH 7.4), 50 mM NaCl, 0.1 mM EDTA,
1 mM DTT, 200 µg/ml BSA, 50% glycerol.

Reaction Conditions:

1X SE-Buffer Y. Incubate at 37° C.

1X SE-Buffer Y (pH 7.9 @ 25° C):

33 mM Tris-Ac 66 mM KAc
10 mM MgAc 1 mM DTT

Heat Inactivation:

Enzyme is inactivated by incubation at 65°C for 20 minutes.

Unit Definition: One unit of the enzyme is the amount required to hydrolyze 1 µg of pUC102 DNA in 1 hour at 37° C in a total reaction volume of 50 µl.

Quality Control Assays

Ligation: After 20-fold overdigestion with Not I, ~90% of the DNA fragments can be ligated and recut.

16-Hour Incubation: A 50 µl reaction containing 1 µg of DNA and 20 Units of enzyme incubated for 16 hours resulted in the same pattern of DNA bands as a reaction incubated for 1 hour.

Oligonucleotide Assay: No detectable degradation of a single-stranded and double-stranded oligonucleotide was observed after incubation with 20 units of restriction endonuclease for 3 hours.

Enzyme Properties:

When using a buffer other than the optimal (Supplied) SE-Buffer, it may be necessary to add more enzymes to achieve complete digestion.

Reagents Supplied with Enzyme:

10X SE Buffer Y.

Blocked by CpG methylation